

COMMENTARY

Childbearing difficulties: A forgotten component of family planning programs in West Africa

DOI: 10.29063/ajrh2022/v26i10.2

Ginette Hounkanrin¹, Rosalie A. Diop², Douaguibé Baguilane³ and Ermel A.K. Johnson^{4,5}*

Women in Global Health, Cotonou, Bénin¹; Institut de population Développement et santé de la Reproduction, Université Cheikh Anta Diop (IPDSR/UCAD), Dakar, Sénégal²; Department Obstetrics and Gynecology, University Teaching Hospital of Lomé and Kara, Kara, Togo³; African Institute for Health Policy & Health Systems, Ebonyi State University, Abakaliki Nigeria⁴; Ecole Nationale des Techniciens en Santé publique et Surveillance Épidémiologique (ENATSE), Université de Parakou, Bénin⁵

*For Correspondence: Email: ermel7@hotmail.com

Introduction

The definition of reproductive health is well codified and agreed upon by all stakeholders: "It's defined as a state of complete physical, mental, and social well-being and not merely the absence of disease or infirmity, in all matters relating to the reproductive system and to its functions and processes. Reproductive health, therefore, implies that people can have a satisfying and safe sex life and that they have the capability to reproduce and the freedom to decide if, when, and how often to do so"¹. This definition makes explicit the ability to decide to reproduce when, how, and how often to do so; implying that every male and female should benefit from interventions to ensure their reproduction.

Historically, family planning programs have evolved from a focus on population control (pre-Cairo) to programs built around four pillars - health, rights, access, and quality of services - after the 1994 Cairo conference¹. While the World Health Organisation (WHO) definition recognizes the two entities of family planning—using contraceptive methods and treating infertility as essential²; more than 25 years after the International Conference on Population and Development (ICPD), the continuing influence of population control in policies and strategies indicates how fertility reduction still motivates much of the programming, research, and advocacy around family planning².

Are the principles of respect for rights and choice reserved exclusively for people who want to

delay having a child or who decide not to have any more children? What about the goal of equity that family planning programs and the new health policy so often advocate? This article examines the silence of family planning programs and interventions of reproductive health on this neglected component of reproductive health and rights services by presenting the extent of reproductive difficulties, their social impact, and the perspectives.

Definition and scope of childbearing difficulties

Infertility is defined by WHO as the inability to conceive a child after 12 months or more of regular unprotected sexual intercourse³⁻⁶. This WHO definition refers to the clinical description of infertility. Beyond this definition, Fortin distinguishes three main types of infertility: (a) female infertility, (b) male infertility, and (c) mixed and unspecified infertility⁷. Whether male or female, infertility can be primary or secondary⁹. In a systematic review, Mascarenhas et al (2012) reported that primary infertility was 1.9% and secondary infertility was 10.5% in women aged 20–44¹⁰. Difficulties in childbearing are observed in both developed and developing countries. Infertility affects 10%-15% of couples worldwide¹¹⁻¹³. According to the WHO, between 50 and 80 million people worldwide suffer from infertility^{14,15}, and trends are not decreasing in Sub-Saharan Africa^{11,16,17} according to studies⁹. The factors of infertility in the couple have female causes (about 50%), male causes (20%-30%), and mixed (20%-

70%)¹¹. Literature reported prevalence of infertility of 9% in The Gambia, 20%-30% in Nigeria¹⁴, 11%-15% in Ghana^{18,19}, 9%-10% in Burkina Faso²⁰. These figures show only a visible part of the magnitude of the problem of reproductive difficulties, which constitutes a public health problem, “a medical disease with a social expression”. Thus, infertility deserves to be put on the agenda, because of its social impact on couples and mostly on women who in the collective imagination have always been historically indexed as the only ones who can suffer from procreation difficulties.

Effects of childbearing difficulties

In Africa, fertility is associated with social recognition or a guarantee of social status. Infertility is a real problem for couples with a significant psychological, social, and economic impact on women^{18,21}. Indeed, procreation is the initial project of marriage in most African cultures. Thus, infertility is more of a social than a medical problem, or rather, a social drama. Regardless of the origin (male or female), and the cause of infertility, the woman most often bears the social burden²². This could be explained by the perception of reproduction in Africa well described by Sow: “African culture makes the woman the vector of reproduction, both biologically and socially. It is in her body that sterility is 'spotted', it is her body that 'betrays' male sterility”²³.

Relationship difficulties and domestic violence, including divorce and remarriage are among the social impacts of infertility on couples' lives^{18,24}. The social control of the family and knowledge contribute to exacerbating the psychological impact with the stigmatization of both the woman and man, thus plunging the affected individuals and couples into a state of permanent distress.

At the individual level, an unfulfilled desire for motherhood has consequences in the different spheres of people's lives. It can be observed that physical and mental health, as well as life plans, are particularly affected by infertility. Feelings and attitudes of anger, especially on the part of the husband, guilt, spite, or grief on the part of the wife, despite all hope, frequently end up jeopardising the meaning and existence of the couple and often end up affecting the household. Most often it is the

arrival of a new wife or simply the divorce. Assisted reproduction treatment for those who can afford them comes with a significant psychological and physical burden, especially for women who undergo most medical procedures. Infertility treatments, in turn, produce feelings of frustration and anger at the medical care received, as well as alternating between the hope of successfully conceiving through medicine and discouragement when treatments fail. Also, gender norms do not make it easy for men to speak out on the subject, particularly as they are struggling with their infertility problem and limit their use of services²⁵. The low use of male reproductive health care by men contributes to an underestimation of the real prevalence and knowledge of associated factors; many men are diagnosed in the context of the couple's search for children²⁶.

Family planning programs are affected by infertility. Indeed, delayed fertility is one of the main barriers to the use of modern contraceptive methods, especially hormonal ones, by young people²⁷. The influence of peers by women with these bad experiences and sociological rumours contributes to the poor results of reproductive health programs.

Family planning programs are silent on reproductive difficulties

Reproductive health, as a reminder, is: “...the general physical, mental and social well-being of the human person, in all matters relating to the reproductive system and to its functions and processes, and not merely the absence of disease or infirmity”¹. This last condition implies that both men and women have the right to access health services that enable women to have a successful pregnancy and birth and give couples every opportunity to have a healthy child. Reproductive health care “the constellation of methods, techniques and services that contribute to reproductive health and well-being by preventing and solving reproductive health problems. It also includes sexual health, the purpose of which is the enhancement of life and personal relations, and not merely counselling and care related to reproduction and sexually transmitted diseases”¹.

Nevertheless, at the country level, infertility interventions are not on the national agenda. In 2008, Sajjabi, the Ugandan presidential

adviser, warned that infertility was not a health priority²⁸. None of the nine countries in the Ouagadougou Partnership has included infertility management interventions as part of family planning activities in their budgeted national action plans²⁹. Similarly, most of the technical partners and donors involved in the field of reproductive health do not propose support or projects for managing infertility cases.

One has the right to wonder about the reasons for this global and collective silence on this aspect of family planning. Many means are made available to women to space out and postpone childbearing, but nothing is offered to them when they face difficulties in procreating at the right time. It is abandoned by all family planning programs, which consider it rather a failure of their interventions. According to Senderowicz, how success is defined, and targets measured on the ground in family planning programs is a clear indication that fertility reduction and contraceptive uptake are, in fact, the primary objectives of these programs, despite the rhetoric about rights and equity². In the absence of more nuanced data, the picture of family planning is commonly summarized by the total fertility rate (TFR), the contraceptive prevalence rate (CPR), and the unmet need for contraceptives to provide a picture of the overall family planning context². However, none of these indicators provides a comprehensive measure of whether the desire for childbearing is being met

Possible solutions for future

Putting infertility on the agenda in reproductive health priorities and programs starts with the availability of research and evaluation data. Generally, a person using the (modern) method is considered a positive outcome, while a person not using the (modern) method is considered a negative outcome². Making infertility effective as an integral part of family planning programs requires, first and foremost, an end to this dichotomous measure of success. It is therefore important to identify new family planning indicators that address problems of difficulty in conceiving, and it is essential that the research community generates information on the current extent of infertility, its psychological and socioeconomic consequences, and possible interventions to support women, couples, and families in seeking care. Recognizing the lack of

answers to many fundamental questions about the prevention, management, and causes of infertility, Duffy *et al.* proposed 10 priorities for future research on male infertility, unexplained female infertility, assisted reproduction, ethics, access, and organization of infertility care³⁰.

Promote and fund interventions that offer solutions to women and couples facing reproductive difficulties. These interventions could include free treatment of sexually transmitted infections and diseases (STIs/STIs), including HIV, the amplification of advocacy for access to infertility care in African countries³¹, and the involvement of the community through civil society organizations. Ombelet and Balen proposed perspectives on infertility, and the integration of infertility in all aspects of reproductive health programs including education, simplification of diagnostic and treatment techniques, and training of health personnel in holistic management³².

Promote and prioritise sexuality education for young people and adolescents. Health education, and specifically the promotion of sexual health among young people/adolescents, has continued to evolve and question the knowledge that young people and adolescents have of their bodies and reproductive systems. This lack of knowledge is a gap and a major handicap for the prevention of infertility.

Strengthening or reorienting health promotion on infertility issues. Indeed, in the context of improving morbidity and mortality, maternal and child health, HIV/AIDS, tuberculosis, malaria and high blood pressure, community health promotion has played a key role in addressing these health problems. A good health promotion campaign can help to raise awareness of the consequences of infertility and contribute to their reduction.

Promote and make accessible modern in vitro fertilisation treatment. The knowledge is now available, even if there are still few centres in black Africa. However, according to Diallo *et al* 1992). "on a continent where sexually transmitted diseases and fertility are on the same upward curve, where social misery and a high birth rate coexist, it may seem paradoxical to talk about assisted reproduction as a 'must' reserved for industrialised countries faced with low birth rates"³³. The social dimension should not be neglected in the search for supportive and caring interventions.

Similarly, interventions should be holistic and include men, given the strong contribution of male factors in couples' infertility⁹. Indeed, psychosocial support by health personnel or social support from the family contributes to the better mental health of infertile couples, without any proven effect on the return of fertility³⁴. In the West African context, social support is important and indispensable as mystical and spiritual factors are put forward as explanations for infertility, contributing to increased stress in couples and recourse to desperate solutions³⁵.

Conclusion

The concept and definition of reproductive health are comprehensive and inclusive, but its translation into interventions obscures an equally important component in the lives of women, couples, and communities: reproductive difficulties. Infertility is a serious societal and public health problem, ignored in reproductive health programs by both countries and technical partners. Because of its impact on the physical, mental, and economic health of the couple and particularly the woman, prevention and management of infertility must be imperative for human rights, equity, and social justice. Country policymakers, reproductive health actors and partners should complement their reproductive health programs with interventions to support women experiencing reproductive difficulties. Thus, the integration of infertility into reproductive health and rights programs, accessibility to diagnosis and reproductive assistance technology should be promoted to ensure women's access to quality infertility care as well as modern contraceptive methods.

Summary box

- Family planning programs have moved from a population control focus (pre-Cairo) to programs built around four pillars: health, rights, access, and quality of services. Childbearing difficulties constitute a real reproductive public health problem in West African countries, given their impact on the mental, physical, economic, and social health of couples.
- Although included in the concept of reproductive health, childbearing difficulties

are not the subject of specific interventions in family planning programs, either at the country level or by most partners.

- Specific interventions, indicators and research on childbearing difficulties should be developed and promotion as part of the family planning programs.

References

1. UNFPA. Programme of action of the international conference on population development. 20th Anniversary Edition. 2014. 296p. ISBN 978-0-89714-022-5 Accessed on 23 June 2022. Available on :https://www.unfpa.org/sites/default/files/pub-pdf/programme_of_action_Web%20ENGLISH.pdf.
2. World Health Organization Department of Reproductive Health and Research (WHO/RHR) and Johns Hopkins Bloomberg School of Public Health/Center for Communication Programs (CCP), Knowledge for Health Project. Family Planning: A Global Handbook for Providers (2018 update). Baltimore and Geneva: CCP and WHO, 2018.
3. Senderowicz L. Contraceptive autonomy: conceptions and measurement of a novel family planning indicator. *Studies in Family Planning*. June 2020. Accessed 6 sept 2022; 51(2):161-76. Available on: <https://onlinelibrary.wiley.com/doi/10.1111/sifp.12114>
4. WHO. Multiple definitions of infertility. Accessed 25 July 2022. Available on: <https://www.who.int/news/item/04-02-2020-multiple-definitions-of-infertility>
5. Zegers-Hochschild F, Adamson GD, de Mouzon J, Ishihara O, Mansour R, Nygren KE, Sullivan S, Vanderpoel, ICMART and WHO. International committee for monitoring assisted reproductive technology (Icmart) and the world health organization (WHO) revised glossary of art terminology, 2009*. *Fertility and Sterility*. nov 2009; 92(5):1520-4. Available on: <https://linkinghub.elsevier.com/retrieve/pii/S0015028209036887>
6. Peterson BD, Newton CR and Rosen KH. Examining congruence between partners' perceived infertility-related stress and its relationship to marital adjustment and depression in infertile couples. *Family Process*. mars 2003; 42(1):59-70. Available on: <https://onlinelibrary.wiley.com/doi/10.1111/j.1545-5300.2003.00059.x>
7. Watkins KJ and Baldo TD. The infertility experience: biopsychosocial effects and suggestions for counselors. *Journal of Counseling & Development*. Oct 2004; 82(4):394-402. Available on: <https://onlinelibrary.wiley.com/doi/10.1002/j.1556-6678.2004.tb00326.x>
8. Fortin, I. Le vécu des hommes et des femmes souffrant ou ayant souffert de difficultés de procréation au Saguenay-Lac-Saint-Jean. Université du Québec à Chicoutimi, 2020. constellation.uqac.ca. <https://constellation.uqac.ca/5748/>.

9. WHO. World Health Organization, *Infertility: A Tabulation of Available Data on Prevalence of Primary and Secondary Infertility*, 1991 Geneva, WHO, WHO/MCH/91.9 accessed on: https://apps.who.int/iris/bitstream/handle/10665/59769/WHO_MCH_91.9.pdf
10. Mascarenhas MN, Flaxman SR, Boerma T, Vanderpoel S and Stevens GA. National, regional, and global trends in infertility prevalence since 1990: a systematic analysis of 277 health surveys. Low N, éditeur. *PLoS Med.* 18 déc 2012; 9(12):e1001356. Available on: <https://dx.plos.org/10.1371/journal.pmed.1001356>
11. Babakhanzadeh E, Nazari M, Ghasemifar S, and Khodadadian A. Some of the Factors Involved in Male Infertility: A Prospective Review. *Int J Gen Med.* 2020 Feb 5;13:29-41. doi: 10.2147/IJGM.S241099. PMID: 32104049; PMCID: PMC7008178.
12. ESHRE Capri Workshop Group, Albertini DF, Anderson R, Bhattacharya S, Evers JLH and McLernon DJ. A prognosis-based approach to infertility: understanding the role of time. *Human Reproduction.* 1 août 2017;32(8):1556-9. Available on: <http://academic.oup.com/humrep/article/32/8/1556/3867334/A-prognosisbased-approach-to-infertility>
13. Fernando Zegers-Hochschild F, Adamson GD, Dyer S, Racowsky C, de Mouzon J, Sokol R, Rienzi L, Sunde A, Schmidt L, Cooke ID, Simpson JL and van der Poel S, The International Glossary on Infertility and Fertility Care, 2017, *Human Reproduction*, Volume 32, Issue 9, September 2017, Pages 1786–1801, <https://doi.org/10.1093/humrep/dex234>
14. Briceag I, Costache A, Purcarea VL, Cergan R, Dumitru M, Briceag I, Sajin M and Ispas AT. Fallopian tubes--literature review of anatomy and etiology in female infertility. *J Med Life.* 2015 Apr-Jun;8(2):129-31. PMID: 25866566; PMCID: PMC4392087.
15. Kumar N and Singh AK. Trends of male factor infertility, an important cause of infertility: a review of literature. *J Hum Reprod Sci.* 2015;8(4):191. doi: 10.4103/0974-1208.170370
16. Ikechebelu JI, Adinma JI, Orie EF and Ikegwuonu SO. High prevalence of male infertility in southeastern Nigeria. *J Obstet Gynaecol* 2003; 23(6): 657-9.
17. Abebe, M.S., Afework, M. & Abaynew, Y. Primary and secondary infertility in Africa: systematic review with meta-analysis. *Fertil Res and Pract* 6, 20 (2020). <https://doi.org/10.1186/s40738-020-00090-3>
18. Chimbatata NBW and Malimba C. Infertility in sub-saharan africa: a woman's issue for how long? A qualitative review of literature. *Open Journal of Social Sciences.* 2016; 04(08):96-102. Available on: <http://www.scirp.org/journal/doi.aspx?doi=10.4236/jss.2016.48012>
19. Asemota O and Klatsky P. Access to infertility care in the developing world: the family promotion gap. *Semin Reprod Med.* 7 janv 2015; 33(01):017-22 <http://dx.doi.org/10.1055/s-0034-1395274>
20. Eric SN, Justine B, Jean NP. Prevalence of the infertility among couples in ouagadougou (Burkina faso): a population-based survey. *TOPHJ [Internet].* 10 nov 2016 [cité 26 oct 2022];9(1):88-97. Disponible sur: <https://openpublichealthjournal.com/VOLUME/9/PAGE/88/>
21. Lampiao F. "It is time the masses are sensitised that men too, like women, have reproductive problems" Fanuel Lampiao talks to Thengo Kavinya on his career in Spermatology. *Malawi Med J.* 2013 Sep;25(3):94. PMID: 24358430; PMCID: PMC3859999.
22. Gulzar, U., Randhawa, R K. and Chaudhary, P. Infertility as a Burden-Women as Victim. *INFERTILITY*, 2021, vol. 7, no 12.
23. Sow F. Stérilité et sous fécondité en Afrique : un drame social. *Revue de Recherche en Santé de la Reproduction en Afrique* P2, ISSN08509700, 1992.
24. Fledderjohann JJ. 'Zero is not good for me': implications of infertility in Ghana, *Human Reproduction*, Volume 27, Issue 5, May 2012, Pages 1383–1390, <https://doi.org/10.1093/humrep/des035>
25. Hanna E and Gough B. The social construction of male infertility: a qualitative questionnaire study of men with a male factor infertility diagnosis. *Sociol Health Illn.* mars 2020;42(3):465-80. Available on: <https://onlinelibrary.wiley.com/doi/10.1111/1467-9566.13038>
26. Mehta A, Nangia AK, Dupree JM and Smith JF. Limitations and barriers in access to care for male factor infertility. *Fertil Steril.* 2016 May;105(5):1128-1137. doi: 10.1016/j.fertnstert.2016.03.023. Epub 2016 Apr 4. PMID: 27054307. doi: 10.1016/j.fertnstert.2016.03
27. Barden-O'Fallon J, Speizer IS, Calhoun LM and Moumouni NA. Return to pregnancy after contraceptive discontinuation to become pregnant: a pooled analysis of West and East African populations. *Reprod Health* 18, 141 (2021). <https://doi.org/10.1186/s12978-021-01193-w>
28. Sajjaji AT. Message from the government of Uganda, *ESHRE Monographs*, Volume 2008, Issue 1, July 2008, Pages 113–114, <https://doi.org/10.1093/humrep/den173>
29. Partenariat de Ouagadougou. Plan d'action budgétisé des pays membres. Accessed at: 23 may 2022. Available on: <https://partenariatouaga.org/plan-daction-des-paysmembres/>.
30. Duffy JMN, Adamson GD, Benson E, Bhattacharya S, Bhattacharya S, Bofill M, Brian K, Collura B, Curtis C, Evers JLH, Farquharson RG, Fincham A, Franik S, Giudice LC, Glanville E, Hickey M, Horne AW, Hull, M. L., Johnson, N. P., V. Jordan, V., Khalaf Y., Knijnenburg, J.M.L., Legro, R.S., Lensen, S., MacKenzie J, Mavrelou D, Mol BW, Morbeck DE, Nagels H, Ng EHY, Niederberger C, Otter AS, Puscasiu, L, Rautakallio-Hokkanen S, Sadler L, Sarris I, Showell M, Stewart JA, Strandell Strawbridge C, Vail A, van Wely M, Vercoe M, Vuong NL, Wang AY, Wang R, Wilkinson J, Wong K, Wong TY, Farquhar CM, and the Priority Setting Partnership for Infertility. Top 10 priorities for future

- infertility research: an international consensus development study. *Fertility and Sterility*. janv 2021; 115(1):180-90.
<https://doi.org/10.1016/j.fertnstert.2020.11.014>.
31. van der Poel SZ. Historical walk: the hrp special programme and infertility. *Gynecol Obstet Invest*. 2012;74(3):218-27. Disponible sur: <https://www.karger.com/Article/FullText/343058>.
32. Ombelet W and van Balen F. Future perspectives. In: Biomedical infertility care in poor resource countries Barriers, Access and Ethics - *Facts, views & vision in ObGyn*. 2012; monograph 87-90 ISSN: 2684-4230 Available on: <https://fvvo.eu/monographs/biomedical-infertility-care-in-poor-resource-countries-barriers-access-and-ethics/>. Accessed: 6 sept 2022.
33. Diallo Y and Biab El Hadi A. Fécondation in vitro en Afrique. *Revue de Recherche en Santé de la Reproduction en Afrique* P21 , ISSN08509700, 1992.
34. Boivin J. A review of psychosocial interventions in infertility. *Social Science & Medicine*. déc 2003; 57(12):2325-41. Available at: <https://linkinghub.elsevier.com/retrieve/pii/S0277953603001382>
35. Tabong PTN and Adongo PB. Understanding the social meaning of infertility and childbearing: a qualitative study of the perception of childbearing and childlessness in northern ghana. Wainberg M, éditeur. *PLoS ONE*. 16 janv 2013; 8(1):e54429. <https://dx.plos.org/10.1371/journal.pone.0054429>.