

ORIGINAL RESEARCH ARTICLE

The effects of caffeine ingestion on blood pressure levels in athletic and non-athletic women

DOI: 10.29063/ajrh2024/v28i5.10

Daniyah K. Al-Shebel and Noorah S Al-Sowayan*

Department of Biology, College of Science, Qassim University, Buraydah, Saudi Arabia

*For Correspondence: Email: nsaoiean@qu.edu.sa

Abstract

Caffeine is one of the most widely consumed pharmacological substances globally, and is known for its potential ergogenic effects. This study examined the impact of caffeine on the blood pressure in athletic and non-athletic women. Caffeine, a CNS stimulant, enhances athletic performance by boosting stamina, alertness, and cognitive speed. The aim of this study was to assess the impact of caffeine on heart rate and blood pressure in both athletic and non-athletic women, and to inform both groups about its effects. The study was conducted in the Kingdom of Saudi Arabia and involved 30 volunteers aged 18–30 years. Participants were equally divided into three groups: athletes who consumed caffeine, non-athletes who consumed caffeine, and a control group (given a placebo). After caffeine ingestion, there were no significant differences in diastolic blood pressure (DBP), systolic blood pressure (SBP), or heart rate between athletes and non-athletes. These findings suggest that caffeine consumption does not significantly affect blood pressure in either athletic or non-athletic women. However, if it raises blood pressure in both groups, it could pose risks, prompting athletes to consider alternative hydration options such as Gatorade. (*Afr J Reprod Health* 2024; 28 [5]: 84-89)

Keywords: Hypertension, systolic blood pressure, diastolic blood pressure, heart rate, coffee, athletes

Résumé

La caféine est l'une des substances pharmacologiques les plus largement consommées dans le monde, et est connue pour ses effets ergogéniques potentiels. Cette étude a examiné l'impact de la caféine sur la pression artérielle des femmes athlètes et non athlètes. La caféine, un stimulant du système nerveux central, améliore les performances des athlètes en augmentant l'endurance, la vigilance et la vitesse cognitive. L'objectif de cette étude était d'évaluer l'impact de la caféine sur la fréquence cardiaque et la pression artérielle chez les femmes athlètes et non athlètes, et d'informer les deux groupes de ses effets. L'étude a été menée au Royaume d'Arabie saoudite et a impliqué 30 volontaires âgés de 18 à 30 ans. Les participants ont été répartis également en trois groupes : des athlètes qui ont consommé de la caféine, des non-athlètes qui ont consommé de la caféine, et un groupe témoin (ayant reçu un placebo). Après l'ingestion de caféine, il n'y avait pas de différences significatives dans la pression artérielle diastolique (PAD), la pression artérielle systolique (PAS) ou la fréquence cardiaque entre les athlètes et les non-athlètes. Ces résultats suggèrent que la consommation de caféine n'affecte pas significativement la pression artérielle chez les femmes, qu'elles soient athlètes ou non. Cependant, si elle augmente la pression artérielle dans les deux groupes, cela pourrait présenter des risques, incitant les athlètes à envisager des options d'hydratation alternatives, telles que le Gatorade. (*Afr J Reprod Health* 2024; 28 [5]: 84-89).

Mots-clés: Hypertension, pression artérielle systolique, pression artérielle diastolique, fréquence cardiaque, café, athlètes

Introduction

Caffeine is the most commonly used pharmacological substance¹. Food sources include coffee, tea, soft drinks, chocolates, energy drinks, and medications²⁻⁴. Coffee contains the highest number of caffeinated products³. There was 65-120 mg caffeine in 230 ml of coffee¹. Healthy people can drink four cups of coffee daily (400 mg of caffeine)¹. Recent phytochemical research has shown the benefits of coffee consumption to human health, as its benefits outweigh its negative effects⁵. Coffee

contains components that may contribute to its vital benefits, such as antioxidant substances, including phenol chlorogenic acid (CA) and tocopherols (TCP)⁶. In recent years, coffee consumption has been associated with a reduced risk of chronic diseases including type 2 diabetes^{10,27}, Parkinson's disease⁹, hepatocellular carcinoma¹¹, endometrial cancer³², and Alzheimer's disease¹².

Coffee consumption is associated with a lower frequency of liver disease and blocked toxin-induced cirrhosis¹. Coffee consumption has a positive protective effect on liver histology and

enzyme levels in patients with non-alcoholic fatty liver disease (NAFLD) and non-alcoholic steatohepatitis (NASH)⁷. One study found that the consumption of two cups of coffee per day decreased the risk of liver cancer by 43%⁸. Clinical research has shown that single doses of caffeine <450 mg do not increase the severity of arrhythmia in healthy individuals⁵.

Elevated caffeine consumption does not increase the risk of incident atrial fibrillation (AF) in healthy women. In contrast, low caffeine consumption may protect against AF²⁸. According to the International Olympic Committee, caffeine positively affects performance¹⁸. Caffeine is a central nervous system stimulant that is now being used to enhance performance in sports³⁸. Athletes are now using caffeine supplements to improve their workouts and performances in competitions³⁹. Moderate doses of caffeine provide an effective ergogenic aid during endurance-type exercise and sports^{29,30}.

Caffeine has been found to boost stamina, and alertness, quicken the speed it takes to process information and logic, and also reduce the amount of effort perceived to be done in an intense exercise period – in doing this, the athlete will feel less fatigued^{38,40}. In addition, caffeine enhances endurance in some activities that last for more than 30 min, possibly because it stimulates the secretion of adrenaline¹⁹ or delays fatigue within the nervous system by blocking or inhibiting adenosine receptors, thus reversing the inhibitory effects of adenosine, such as decreasing the synthesis of DA (a neurotransmitter) associated with an excess of 5-HT, and thus delaying fatigue¹⁷.

In a previous study, caffeine did not increase SBP or DBP when maximal exercise was performed in physically active females¹³. Another study found that caffeine consumption (4 mg/kg) did not affect the cardiovascular system after exercise¹⁴. In two studies, caffeine consumption significantly increased Post-exercise SBP^{15,16}. It is anticipated that 29% of adults living in developing nations (1.56 billion) will have hypertension by 2025²¹. Therefore, to avoid and reduce the risk of morbidity and mortality due to cardiovascular disease, blood pressure monitoring is necessary to maintain appropriate blood pressure levels⁴⁴. Therefore, digital blood pressure monitors have received increasing attention as alternatives to aneroid and

mercury manometers because they require qualified healthcare workers to use them⁴⁵. Some studies recommend the use of digital blood pressure monitors^{46,47}.

In one study, blood pressure increased after caffeine ingestion (200-300 mg) in hypertensive patients 60 min after ingestion²⁰. Additionally, they discovered in another study that healthy older normotensive women who consumed caffeine experienced a substantial rise in systolic and diastolic blood pressure (DBP) that persisted for 90 minutes³¹. In another study, they found that in a hypertensive older population, habitual coffee ingestion of >3 cups/day was associated with uncontrolled BP²². In two other studies, neither caffeinated nor decaffeinated coffee was linked to higher blood pressure in the long term^{23,24}. Coffee consumers have a higher risk of developing hypertension than non-coffee drinkers²⁴. The effect of caffeine on blood pressure is due to its ability to inhibit adenosine receptors in the blood vessels, which causing vasoconstriction^{25,26}. The level of physical activity and variation in DBP after caffeine ingestion were inversely correlated in one study, indicating that younger women who engaged in less physical activity experienced an increase in DBP after caffeine ingestion^{31,33}.

The purpose of our study was to determine the effects of caffeine consumption on both athletic and non-athletic females. We wanted to determine whether caffeine raised blood pressure in both types and determine if it had a greater effect on one, so that we could inform the population about caffeine in their lifestyle.

If it was known that caffeine raised high blood pressure in a non-trained individual, and at the same time, the same effects were shown in an athlete, it could be considered dangerous for the non-athlete. This would be the same if caffeine had no effect on the athlete at all, and the athlete may want to reconsider consuming the product, considering that there are no effects from the caffeine shown on the blood pressure, and would be better off with regular Gatorade⁴¹⁻⁴³.

Methods

This study was conducted on 30 volunteers with ages ranging 18-30 years. Participants with hypertension or cardiovascular disease were excluded. Participants were requested to avoid consuming any

source of caffeine, including medications, one day before the experiment (including medications).

The participants were divided equally into 3 groups:

(i)- Athletes who ingested caffeine

(ii) Non-athletes (normal) who ingested caffeine.

(iii) Control (placebo)

Information such as (weight - height - range of daily caffeine intake) was recorded for all participants before the experiment began. Groups (i) and (ii) ingested 250-300 ml of coffee (average 80–120 mg of caffeine), and group (iii) ingested 250-300 ml of water. SBP, DBP, and heart rate were measured for all groups before and after ingestion (–45-90 minutes after ingestion because within 45 min after oral ingestion, caffeine is completely absorbed in the small intestine and stomach; it reaches its highest concentration in the plasma within 20-30 min after ingestion¹. Measurements were performed in a left-handed resting position using an automatic “digital” blood pressure monitor (CITIZEN, CH-657).

Results

Analysis of variance (ANOVA) was conducted using the SPSS software. Data were evaluated using descriptive statistical analyses and paired t-tests. All data processing was done at a probability value $p < 0.05$. The average age of participants was 22.40 ± 3.41 and weight was 59.93 ± 9.24 .

Table 1: Demographic data for participants' included standard deviation and average values

	N	Mean	Std. Deviation
Age	30	22.40	3.41
Weight	30	59.93	9.24
Height	30	156.03	5.17
No. of coffee cups per day	30	1.27	0.98
Valid N (listwise)	30		

Table 1 shows the demographic data of the participants, including standard deviation and average values. Table 2 displays the findings following the use of the paired t-test to compare variations in SBP, DBP, and heart rate between athletes, non-athletes, and controls after ingestion of caffeine and placebo. The results of the paired samples test are summarized in Table 3 as the mean values of SBP, DBP, and Pulse in athletes and non-

Table 2: The standard deviation and mean of the variables are shown in the overview of the paired t-test between athletes, non-athletes, and control

	N	Mean	Std. Deviation
VAR00010			
SYS_ Normal (After)	10	113.30	6.83
SYS_ athletes (After)	10	116.10	11.43
SYS_ control (After)	10	116.70	8.62
DIA_ Normal (After)	10	83.30	8.03
DIA_ athletes (After)	10	85.80	9.27
DIA_ control (After)	10	84.60	7.62
Pulse_ Normal (After)	10	98.90	13.44
Pulse_ athletes (After)	10	86.80	17.54
Pulse_ control (After)	10	86.70	6.70

athletes after caffeine ingestion and in the control group.

The average SBP difference of non-athletes after drinking a coffee was 2.80 ± 5.20 ($t = 1.702$, $p = 0.123 > 0.05$), and of athletes it was -2.5 ± 10.92 ($t = -0.724$, $p = 0.487 > 0.05$). The average DBP variation or difference of non-athletes after consuming coffee was 1.30 ± 6.07 ($t = 0.677$, $p = 0.516 > 0.05$) and of athletes it was -4.7 ± 7.2 ($t = -2.065$, $p = 0.069 > 0.05$). The typical heart rate difference of non-athletes after consumption of coffee was -1.90 ± 13.55 ($t = -0.443$, $p = 0.668 > 0.05$), and that of athletes was 0.8 ± 12.32 ($t = 0.205$, $p = 0.842 > 0.05$). Consequently, we concluded that there was no difference in DBP, SBP, or heart rate between athletes and non-athletes after caffeine ingestion. (Table 3) There was a slight increase in SBP and DBP in athletic women after caffeine consumption.

Discussion

In our study, we found that caffeine ingestion had no effect on blood pressure in both athletic and non-athletic women. This result was supported by some studies, in two of them they found that moderate or habitual caffeine intake (> 3 cups/day or 440 mg of caffeine/day) was not associated with the risk of hypertension^{36,34}. A systematic review found that moderate caffeine consumption was inversely associated with the risk of cardiovascular disease (CVD), whereas high coffee consumption was not associated with a higher risk of CVD³⁷. A meta-analysis of 21 prospective cohort studies found that moderate caffeine consumption was associated with a lower risk of coronary heart disease (CHD) in women³⁵. Our study has some limitations.

Table 3: BP and heart rates test results

Paired Differences and Samples Test	Mean	Std. Deviation	t	df	Sig	
Samples of Normal (with caffeine)	SYS (Before) -SYS (After)	2.80	5.20	1.702	9	.123
	DIA (Before) - DIA (After)	1.30	6.07	.677	9	.516
	Pulse (Before) - Pulse (After)	-1.90	13.55	-.443	9	.668
Samples of athletes (with caffeine)	SYS (Before) -SYS (After)	-2.5	10.92	-.724	9	.487
	DIA (Before) - DIA (After)	-4.7	7.2	-2.065	9	.069
	Pulse (Before) - Pulse (After)	0.8	12.32	.205	9	.842
Control	SYS (Before) -SYS (After)	0.7	6.11	.362	9	.726
	DIA (Before) - DIA (After)	1.3	4.62	.890	9	.397
	Pulse (Before) - Pulse (After)	1.4	9.63	.460	9	.657

The sample size was small, may be considered non-representative, and the study time was short. We recommend further investigation of the impact of caffeine on human health and blood pressure in long-term large-scale studies.

In summary, the results of this study indicate that there is no relationship between caffeine ingestion and blood pressure in both athletic and non-athletic women. However, if it raises blood pressure in both groups, it could pose risks, prompting athletes to consider alternative hydration options such as Gatorade.

Conclusion

Author contributions

Daniyah Al-Shebel performed the investigation, resources, methodology, validation, data curation, formal analysis, and preparation of the original draft. Al-Shebel conducted the resources, data curation, and formal analysis.

Noorah Al-Sowayan conceptualized the methodology, validation, writing—review and editing, visualization, supervision, and project administration.

All authors have read and agreed to the published version of the manuscript.

Consent form

A meeting was held with the volunteers and the experiment was explained to them. We obtained their consent and signed an informed consent form. They were informed that they could leave the experiment at any time.

Funding

This study received no external funding.

Data availability statement

Data is contained within the article.

Conflicts of interest

The authors declare no conflicts of interest.

References

1. Nieber K. The Impact of Coffee on Health. *Planta Med.* 2017 Nov;83(16):1256-1263. doi: 10.1055/s-0043-115007. Epub 2017 Jul 4.
2. Mitchell DC, Knight CA, Hockenberry J, Teplansky R and Hartman TJ. Beverage caffeine intakes in the U.S. *Food Chem Toxicol.* 2014 Jan;63:136-42. doi: 10.1016/j.fct.2013.10.042. Epub 2013 Nov 1. PMID: 24189158.
3. Reyes CM and Cornelis MC. Caffeine in the Diet: Country-Level Consumption and Guidelines. *Nutrients.* 2018;10(11):1772. Published 2018 Nov 15. doi:10.3390/nu10111772
4. Heckman MA, Weil J and Gonzalez de Mejia E. Caffeine (1, 3, 7-trimethylxanthine) in foods: a comprehensive review on consumption, functionality, safety, and regulatory matters. *J Food Sci.* 2010 Apr;75(3):R77-87. doi: 10.1111/j.1750-3841.2010.01561.x. PMID: 20492310.
5. Nawrot P, Jordan S, Eastwood J, Rotstein J and Hugenholtz A. and Feeley M. Effects of caffeine on human health. *Food Addit Contam.* 2003 Jan;20(1):1-30. doi: 10.1080/0265203021000007840.
6. Johnston KL, Clifford MN and Morgan LM. Coffee acutely modifies gastrointestinal hormone secretion and glucose tolerance in humans: glycemic effects of chlorogenic acid and caffeine. *Am J Clin Nutr.* 2003 Oct;78(4):728-33. doi: 10.1093/ajcn/78.4.728.

7. Degertekin B, Tozun N, Soylemez AG, Gurtay E, Bozkurt U, Yilmaz Y, Yapali S, Vardareli E, Unal HU, Colakoglu B and Alpaydin CB. Regular coffee intake improves liver enzyme levels and liver histology in patients with chronic alcohol consumption, non-alcoholic fatty liver and non-alcoholic steatohepatitis: Report of 259 cases. *Hepatol Forum.* 2020;1(3):88-96. Published 2020 Sep 21. doi:10.14744/hf.2020.2020.0026
8. Larsson SC and Wolk A. Coffee consumption and risk of liver cancer: a meta-analysis. *Gastroenterology.* 2007 May;132(5):1740-5. doi: 10.1053/j.gastro.2007.03.044. Epub 2007 Mar 24. PMID: 17484871.
9. Ascherio A, Zhang SM, Hernán MA, Kawachi I, Colditz GA, Speizer FE and Willett WC. Prospective study of caffeine consumption and risk of Parkinson's disease in men and women. *Ann Neurol.* 2001 Jul;50(1):56-63. doi: 10.1002/ana.1052.
10. Salazar-Martinez E, Willett WC, Ascherio A, Manson JE, Leitzmann MF, Stampfer MJ and Hu FB. Coffee consumption and risk for type 2 diabetes mellitus. *Ann Intern Med.* 2004 Jan 6;140(1):1-8. doi: 10.7326/0003-4819-140-1-200401060-00005.
11. Inoue M, Yoshimi I, Sobue T, Tsugane S and JPHC Study Group. Influence of coffee drinking on subsequent risk of hepatocellular carcinoma: a prospective study in Japan. *J Natl Cancer Inst.* 2005 Feb 16;97(4):293-300. doi: 10.1093/jnci/dji040.
12. Arendash GW and Cao C. Caffeine and coffee as therapeutics against Alzheimer's disease. *J Alzheimers Dis.* 2010;20 Suppl 1:S117-26. doi: 10.3233/JAD-2010-091249.
13. Connahan LE, Ott CA and Barry VW. Effect of Caffeine on near Maximal Blood Pressure and Blood Pressure Recovery in Physically-Active, College-Aged Females. *Int J Exerc Sci.* 2017;10(2):266-273. Published 2017 Mar 1.
14. Joao Bruno YC, Anunciacao PG, Ruiz RJ, Polito MD, and Casonatto J. Effect of caffeine intake on blood pressure and heart rate variability after a single bout of aerobic exercise : original research article. *International SportMed Journal* Vol. 13, No. 3. 2012. doi:abs/10.10520/EJC125879.
15. Goldstein E, Jacobs PL, Whitehurst M, Penhollow T and Antonio J. Caffeine enhances upper body strength in resistance-trained women. *J Int Soc Sports Nutr.* 2010;7:18. Published 2010 May 14. doi:10.1186/1550-2783-7-18
16. Woolf K, Bidwell WK and Carlson AG. The effect of caffeine as an ergogenic aid in anaerobic exercise. *Int J Sport Nutr Exerc Metab.* 2008 Aug;18(4):412-29. doi: 10.1123/ijnsnem.18.4.412. PMID: 18708685.
17. Davis JM, Zhao Z, Stock HS, Mehl KA, Buggy J and Hand GA. Central nervous system effects of caffeine and adenosine on fatigue. *Am J Physiol Regul Integr Comp Physiol.* 2003 Feb;284(2):R399-404. doi: 10.1152/ajpregu.00386.2002. Epub 2002 Oct 24.
18. Maughan RJ, Burke LM, Dvorak J, Larson-Meyer DE, Peeling P, Phillips SM, Rawson ES, Walsh NP, Garthe I, Geyer H, Meeusen R, van Loon LJC, Shirreffs SM, Spriet LL, Stuart M, Vernec A, Currell K, Ali VM, Budgett RG, Ljungqvist A, Mountjoy M, Pitsiladis YP, Soligard T, Erdener U and Engebretsen L. IOC consensus statement: dietary supplements and the high-performance athlete. *Br J Sports Med.* 2018 Apr;52(7):439-455. doi: 10.1136/bjsports-2018-099027. Epub 2018 Mar 14. PMID: 29540367.
19. Graham TE. Caffeine and exercise: metabolism, endurance and performance. *Sports Med.* 2001;31(11):785-807. doi: 10.2165/00007256-200131110-00002.
20. Mesas AE, Leon-Muñoz LM, Rodriguez-Artalejo F and Lopez-Garcia E. The effect of coffee on blood pressure and cardiovascular disease in hypertensive individuals: a systematic review and meta-analysis. *Am J Clin Nutr.* 2011 Oct;94(4):1113-26. doi: 10.3945/ajcn.111.016667. Epub 2011 Aug 31.
21. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK and He J. Global burden of hypertension: analysis of worldwide data. *Lancet.* 2005 Jan 15-21;365(9455):217-23. doi: 10.1016/S0140-6736(05)17741-1.
22. Lopez-Garcia E, Orozco-Arbeláez E, Leon-Muñoz LM, Guallar-Castillon P, Graciani A, Banegas JR and Rodríguez-Artalejo F. Habitual coffee consumption and 24-h blood pressure control in older adults with hypertension. *Clin Nutr.* 2016 Dec;35(6):1457-1463. doi: 10.1016/j.clnu.2016.03.021.
23. Winkelmayr WC, Stampfer MJ, Willett WC and Curhan GC. Habitual caffeine intake and the risk of hypertension in women. *JAMA.* 2005 Nov 9;294(18):2330-5. doi: 10.1001/jama.294.18.2330.
24. Klag MJ, Wang NY, Meoni LA, Brancati FL, Cooper LA, Liang KY, Young JH and Ford DE. Coffee intake and risk of hypertension: the Johns Hopkins precursors study. *Arch Intern Med.* 2002 Mar 25;162(6):657-62. doi: 10.1001/archinte.162.6.657.
25. Umemura T, Ueda K, Nishioka K, Hidaka T, Takemoto H, Nakamura S, Jitsuiki D, Soga J, Goto C, Chayama K, Yoshizumi M and Higashi Y. Effects of acute administration of caffeine on vascular function. *Am J Cardiol.* 2006 Dec 1;98(11):1538-41. doi: 10.1016/j.amjcard.2006.06.058.
26. Ozkan B, Yüksel N, Anik Y, Altintas O, Demirci A and Çağlar Y. The effect of caffeine on retrolubar hemodynamics. *Curr Eye Res.* 2008 Sep;33(9):804-9. doi: 10.1080/02713680802344708.
27. van Dam RM and Feskens EJ. Coffee consumption and risk of type 2 diabetes mellitus. *Lancet.* 2002 Nov 9;360(9344):1477-8. doi: 10.1016/S0140-6736(02)11436-X.
28. Conen D, Chiuvè SE, Everett BM, Zhang SM, Buring JE and Albert CM. Caffeine consumption and incident atrial fibrillation in women. *Am J Clin Nutr.* 2010 Sep;92(3):509-14. doi: 10.3945/ajcn.2010.29627. Epub 2010 Jun 23. PMID: 20573799.
29. Guest NS, VanDusseldorp TA, Nelson MT, Grgic J, Schoenfeld BJ, Jenkins NDM, Arent SM, Antonio J, Stout JR, Trexler ET, Smith-Ryan AE, Goldstein ER, Kalman DS and Campbell BI. International society of sports nutrition position stand: caffeine and exercise performance. *J Int Soc Sports Nutr.* 2021 Jan 2;18(1):1. doi: 10.1186/s12970-020-00383-4.
30. Lima-Silva AE, Cristina-Souza G, Silva-Cavalcante MD, Bertuzzi R and Bishop DJ. Caffeine during High-

- Intensity Whole-Body Exercise: An Integrative Approach beyond the Central Nervous System. *Nutrients*. 2021;13(8):2503. Published 2021 Jul 22. doi:10.3390/nu13082503.
31. Arciero PJ and Ormsbee MJ. Relationship of blood pressure, behavioral mood state, and physical activity following caffeine ingestion in younger and older women. *Appl Physiol Nutr Metab*. 2009 Aug;34(4):754-62. doi: 10.1139/H09-068.
 32. Je Y and Giovannucci E. Coffee consumption and risk of endometrial cancer: findings from a large up-to-date meta-analysis. *Int J Cancer*. 2012 Oct 1;131(7):1700-10. doi: 10.1002/ijc.27408. Epub 2012 Jan 31.
 33. Zhang Z, Hu G, Caballero B, Appel L and Chen L. Habitual coffee consumption and risk of hypertension: a systematic review and meta-analysis of prospective observational studies. *Am J Clin Nutr*. 2011 Jun;93(6):1212-9. doi: 10.3945/ajcn.110.004044. Epub 2011 Mar 30.
 34. Grosso G, Micek A, Godos J, Sciacca S, Pajak A, Martínez-González MA, Giovannucci EL and Galvano F. Coffee consumption and risk of all-cause, cardiovascular, and cancer mortality in smokers and non-smokers: a dose-response meta-analysis. *Eur J Epidemiol*. 2016 Dec;31(12):1191-1205. doi: 10.1007/s10654-016-0202-2. Epub 2016 Oct 3.
 35. Wu JN, Ho SC, Zhou C, Ling WH, Chen WQ, Wang CL and Chen YM. Coffee consumption and risk of coronary heart diseases: a meta-analysis of 21 prospective cohort studies. *Int J Cardiol*. 2009 Nov 12;137(3):216-25. doi: 10.1016/j.ijcard.2008.06.051. Epub 2008 Aug 15. PMID: 18707777.
 36. Ciaramelli C, Palmioli A and Airoldi C. Coffee variety, origin and extraction procedure: Implications for coffee beneficial effects on human health. *Food Chem*. 2019 Apr 25; 278:47-55. doi: 10.1016/j.foodchem.2018.11.063. Epub 2018 Nov 12.
 37. Mendoza MF, Sulague RM, Posas-Mendoza T and Lavie CJ. Impact of Coffee Consumption on Cardiovascular Health. *Ochsner J*. 2023;23(2):152-158. doi:10.31486/toj.22.0073.
 38. Jodra P, Lago-Rodríguez A, Sánchez-Oliver AJ, López-Samanes A, Pérez-López A, Veiga-Herreros P, San Juan AF and Domínguez R. Effects of caffeine supplementation on physical performance and mood dimensions in elite and trained-recreational athletes. *J Int Soc Sports Nutr*. 2020 Jan 3;17(1):2. doi: 10.1186/s12970-019-0332-5.
 39. Jiménez SL, Díaz-Lara J, Pareja-Galeano H and Del Coso J. Caffeinated Drinks and Physical Performance in Sport: A Systematic Review. *Nutrients*. 2021 Aug 25;13(9):2944. doi: 10.3390/nu13092944.
 40. Lopes-Silva JP, Rocha ALS, Rocha JCC, Silva VFDS and Correia-Oliveira CR. Caffeine ingestion increases the upper-body intermittent dynamic strength endurance performance of combat sports athletes. *Eur J Sport Sci*. 2022 Feb;22(2):227-236. doi: 10.1080/17461391.2021.1874058.
 41. Shyamali Kaushalya F, Romero-Arenas S, García-Ramos A, Colomer-Poveda D and Marquez G. Acute effects of transcranial direct current stimulation on cycling and running performance. A systematic review and meta-analysis. *Eur J Sport Sci*. 2022 Feb;22(2):113-125. doi: 10.1080/17461391.2020.1856933.
 42. O'Hara, Reginald B and William W. Dodson. Sublingual Caffeine Supplementation and Its Effects on Physical Performance Measures in Highly Fit United States Military Personnel. *International Journal of Exercise Science: Conference Proceedings*. 2023.
 43. de Oliveira LF, Dolan E, Swinton PA, Durkalec-Michalski K, Artioli GG, McNaughton LR and Saunders B. Extracellular Buffering Supplements to Improve Exercise Capacity and Performance: A Comprehensive Systematic Review and Meta-analysis. *Sports Med*. 2022 Mar;52(3):505-526. doi: 10.1007/s40279-021-01575-x.
 44. Menezes AM, Dumith SC, Noal RB, Nunes AP, Mendonça FI, Araújo CL, Duval MA, Caruso PE and Hallal PC. Validity of a wrist digital monitor for blood pressure measurement in comparison to a mercury sphygmomanometer. *Arq Bras Cardiol*. 2010. doi: 10.1590/S0066-782X2010000300014.
 45. Stergiou GS, Lin CW, Lin CM, Chang SL, Protogerou AD, Tzamouranis D, Nasothimiou E and Tan TM. Automated device that complies with current guidelines for office blood pressure measurement: design and pilot application study of the Microlife WatchBP Office device. *Blood Press Monit*. 2008 Aug;13(4):231-5. doi: 10.1097/MBP.0b013e3283057a84.
 46. Yoo S, Baek H, Doh K, Jeong J, Ahn S, Oh IY and Kim K. Validation of the mobile wireless digital automatic blood pressure monitor using the cuff pressure oscillometric method, for clinical use and self-management, according to international protocols. *Biomed Eng Lett*. 2018 Sep 21;8(4):399-404. doi: 10.1007/s13534-018-0085-0.
 47. Kario K, Shimbo D, Tomitani N, Kanegae H, Schwartz JE and Williams B. The first study comparing a wearable watch-type blood pressure monitor with a conventional ambulatory blood pressure monitor on in-office and out-of-office settings. *J Clin Hypertens (Greenwich)*. 2020 Feb;22(2):135-141. doi: 10.1111/jch.13799.